

## PART III: CONSUMER INFORMATION

Pr **PRO-BACLOFEN**

**Baclofen Tablets USP**

This leaflet is part III of a three-part -"Product Monograph" published when PRO-BACLOFEN was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about PRO-BACLOFEN. Contact your doctor or pharmacist if you have any questions about the drug.

### ABOUT THIS MEDICATION

#### What the medication is used for:

PRO-BACLOFEN is used to reduce and relieve the excessive stiffness and/or spasms in your muscles occurring in various conditions such as multiple sclerosis and diseases or injuries of the spinal cord.

#### What it does:

Due to the relaxation of muscle and the consequent relief from pain, PRO-BACLOFEN improves your ability to move, makes it easier for you to manage your daily activities and facilitates physiotherapy.

If you have any questions about how PRO-BACLOFEN works or why this medicine has been prescribed for you, ask your doctor.

#### When it should not be used:

Do not take PRO-BACLOFEN

- If you are allergic (hypersensitive) to baclofen or any of the other ingredients listed in "**What the nonmedicinal ingredients are**".

If this applies to you, tell your doctor without taking PRO-BACLOFEN.

If you think you may be allergic, ask your doctor for advice.

#### What the medicinal ingredient is

The active substance of PRO-BACLOFEN is baclofen.

#### What the non-medicinal ingredients are

The non-medicinal ingredients are: microcrystalline cellulose, starch, magnesium stearate and lactose.

#### What dosage forms it comes in:

PRO-BACLOFEN is available in 10 mg and 20 mg tablets.

### WARNINGS AND PRECAUTIONS

#### SERIOUS WARNINGS AND PRECAUTIONS

Taking PRO-BACLOFEN with opioid medicines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.

BEFORE you use PRO-BACLOFEN talk to your doctor or pharmacist if you:

- Have kidney disease. Your doctor will decide whether or not PRO-BACLOFEN is the appropriate treatment for you;
- Are suffering from epilepsy (seizures);
- Have acute pain in your stomach (ulcer) or intestines, breathing problems, liver disease, or a disturbance of blood circulation in your brain.
- Are taking medicines for arthritis or pain (see section: "**Interactions with this medication**").
- Have difficulty urinating;
- Have Parkinson's disease or certain mental illnesses accompanied by confusion or depression;
- Are diabetic.
- Have or had thoughts of harming or killing yourself, have or had depression, have a history of alcoholism or drink alcohol to excess, have a history of drug abuse and dependence, or are taking medicines associated with increased suicide risk;
- Have thoughts of harming or killing yourself at any time, speak to your doctor straightaway or go to a hospital. Also, ask a relative or close friend to tell you if they are worried about any changes in your behavior and ask them to read this leaflet

#### **Older people (aged 65 years or above) or people with a disturbance of circulation in the brain**

If you are in one of these groups, you may experience more side effects. Therefore, your doctor will keep you under appropriate surveillance and may adapt the dose of PRO-BACLOFEN you take.

#### **Children and adolescents**

Safe use of baclofen in children under age 12 has not been established and PRO-BACLOFEN is therefore not recommended for use in children.



### Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

You should not use PRO-BACLOFEN during pregnancy unless your doctor advises you to do so. Tell your doctor if you are pregnant, planning to become pregnant, or breast-feeding. He or she will discuss with you the potential risk of taking PRO-BACLOFEN during pregnancy or if you are breast-feeding. Use of PRO-BACLOFEN during pregnancy may result in the newborn experiencing withdrawal from the drug including, irritability, high-pitched crying, trembling, increased muscle tone, excessive sucking, disordered sleep, increase in body temperature, uneven discolored patches on the skin, and convulsions and other symptoms related to sudden stop of treatment sometime after delivery. Your doctor may need to treat your newborn for withdrawal reactions.

### Driving and using machines

In some people, PRO-BACLOFEN may be associated with dizziness, sleepiness or visual disturbance. If this happens to you, do not drive a car, use a machine, or do other things that need your full attention.

### Further safety measures

Before having any kind of surgery (including by the dentist), or emergency treatment, tell the doctor in charge that you are taking PRO-BACLOFEN.

### INTERACTIONS WITH THIS MEDICATION

Drugs that may interact with PRO-BACLOFEN include:

- Alcohol;
- Sedative drugs;
- Medicines used to treat mood disorders such as antidepressants and lithium;
- Medicines used to treat high blood pressure;
- Medicines used to treat Parkinson's disease;
- Medicines for arthritis or pain.

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

You should not drink alcohol during your treatment with PRO-BACLOFEN.

### PROPER USE OF THIS MEDICATION

#### Usual dose:

Follow your doctor's instructions carefully. Do not exceed the recommended dose.

#### **How much PRO-BACLOFEN to take**

Treatment usually starts with 15 mg daily, preferably taken in divided doses. The dose is then gradually increased until the best results are obtained; this may be between 40 mg to 80 mg per day, taken in divided doses.

The dose prescribed by your doctor may be different from that written here. If this is the case, follow the doctor's instructions.

Your doctor will tell you exactly how many tablets of PRO-BACLOFEN to take.

Depending on how you respond to the treatment, your doctor may suggest a higher or lower dose.

#### **When to take PRO-BACLOFEN**

Taking PRO-BACLOFEN at the same time each day will help you to remember when to take your medicine.

#### **How to take PRO-BACLOFEN**

Be sure to take this medicine regularly, and exactly as your doctor tells you. This will help you to get the best results and reduce the risk of side effects.

#### **How long to take PRO-BACLOFEN**

Continue taking PRO-BACLOFEN as your doctor tells you.

If you have questions about how long to take PRO-BACLOFEN talk to your doctor or your pharmacist.

**Do not suddenly stop taking PRO-BACLOFEN without first checking with your doctor.** He or she will tell you when and how you can stop taking this medicine; stopping suddenly can make your condition worse.

**If you stop your treatment suddenly, you may experience:** nervousness, feeling confused, hallucinations, abnormal thinking or behaviour, convulsions, uncontrollable twitching, jerking or writhing movements, fast heart beat, high body temperature. The excessive stiffness (spasms) in your muscles may also worsen.

#### Overdose



If you think you have taken too much PRO-BACLOFEN contact your healthcare professional, hospital emergency department or regional poison control center immediately, even if there are no symptoms

If you have accidentally taken many more tablets than your doctor has prescribed, seek immediate emergency medical treatment, even though you do not feel sick.

The main symptoms of overdose are drowsiness, breathing difficulties, trouble of consciousness and being unconscious (coma).

Other symptoms may include: feeling confused, hallucinations, agitation, convulsions, blurred vision, unusual muscle weakness, sudden contraction of the muscles, poor or absent reflexes, high or low blood pressure, slow, fast or irregular heart beat, low body temperature, nausea, vomiting, diarrhea or excessive salivation, trouble breathing during sleep (sleep apnoea), pain in muscles, fever and dark urine (rhabdomyolysis).

If you have **kidney disease** and have accidentally taken more tablets or more syrup than **your doctor** has prescribed, you may experience neurological symptoms of overdose (e.g. drowsiness, feeling confused, hallucinations).

### Missed Dose

If you have forgotten to take one of your scheduled doses, take it as soon as you remember. However, if it is almost time for your next dose, do not take the missed one at the same time as the scheduled one, otherwise you will be doubling the dose. Just go back to your regular dosing timetable. If you have forgotten to take several doses you should contact your doctor.

### SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, PRO-BACLOFEN can have some side effects, although not everybody gets them. These are often mild and are usually at the start of treatment; they normally wear off after a few days.

#### Very common side effects

*These side effects may affect more than 1 in 10 patients:*

- Drowsiness, sleepiness;
- Nausea.

If any of these affects you severely, **tell your doctor**.

#### Common side effects

*These side effects may affect between 1 and 10 in every 100 patients:*

- Feeling faint, tiredness, dizziness, headache, inability to sleep, weakness in arms and legs, pain in muscles, uncontrollable eye movements, dry mouth;
- Disturbance of the digestive tract, retching, vomiting, constipation, diarrhea;
- Sweating a lot;
- Passing more urine than normal, bedwetting.

If any of these affects you severely, **tell your doctor**.

#### Rare side effects

*These side effects may affect between 1 and 10 in every 10,000 patients:*

- Tingling or numbness of the hands and/or feet, difficulty in speaking, taste disturbance;
- Abdominal pain;
- Sudden decrease in urine;
- Inability to get or to maintain an erection (impotence).

#### Side effect also reported (frequency unknown)

- Increased blood sugar;
- Drug misuse, drug abuse, drug dependence.

If any of these affects you severely, **tell your doctor**.

If you notice any other side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

#### SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom/effect		Talk with your doctor or pharmacist		Stop taking drug and seek immediate emergency medical treatment
		Only if severe	In all cases	
Common	Breathing problems		√	
	Feeling of confusion		√	
	Feeling of extreme happiness		√	
	Sad mood (depression)		√	
	Loss of coordination affecting balance and walking, limb and eye movements and/or speech (signs of ataxia)		√	
	Trembling		√	
	Hallucinations		√	
	Nightmares		√	



	Blurred vision/visual disturbance		√	
	Shortness of breath at rest or with activity, swelling in the legs and tiredness (signs of decreased cardiac output)		√	
	Low blood pressure (hypotension)		√	
	Skin rash and hives		√	
	Difficulty passing urine, pain when passing urine or a sudden decrease in urine		√	
Rare	Abdominal pain, yellowing of the skin or eyes and tiredness (signs of liver disturbance)			√
	Convulsions			√
Very rare	Low body temperature		√	
Not known	Symptoms following sudden discontinuation of the medicine (drug withdrawal syndrome)		√	
	Slow heart beat		√	
	Suicidal Behavior: thoughts or actions about harming or killing yourself		√	

*This is not a complete list of side effects. For any unexpected effects while taking PRO-BACLOFEN, contact your doctor or pharmacist.*

#### HOW TO STORE IT

PRO-BACLOFEN Tablets should be stored in tight containers between 15°C to 30°C.

Keep out of the reach and sight of children.

#### REPORTING SUSPECTED SIDE EFFECTS

##### Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

#### MORE INFORMATION

##### If you want more information about PRO-BACLOFEN:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Consumer Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); or by contacting Pro Doc Ltée at 1-800-361-8559, [www.prodoc.qc.ca](http://www.prodoc.qc.ca) or [info@prodoc.qc.ca](mailto:info@prodoc.qc.ca).

This leaflet was prepared by  
Pro Doc Ltée, Laval, Québec, H7L 3W9

Last revised: APR 17, 2025