

PART III: CONSUMER INFORMATION

Pr PRO-RAMIPRIL – 1,25

Pr PRO-RAMIPRIL – 2,5

Pr PRO-RAMIPRIL – 5

Pr PRO-RAMIPRIL – 10

Ramipril Capsules pro Doc Standard

This leaflet is part III of a three-part “Product Monograph” published when PRO-RAMIPRIL was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about PRO-RAMIPRIL. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

High Blood Pressure (Hypertension)

PRO-RAMIPRIL lowers high blood pressure. It can be used alone or together with a diuretic (“water pill”).

Managing your lifestyle

Keeping your blood pressure controlled

It takes more than just medication to reduce blood pressure. Discuss the risk factors, and how they apply to your lifestyle, with your doctor. You may have to modify some of your daily habits to keep your blood pressure down.

Exercise regularly. It will help to keep your weight down, make you feel more energetic and is a good way to deal with stress. If you are not exercising regularly, be sure to discuss a fitness plan with your doctor.

Remember, hypertension is a long-term disease without symptoms. Just because you feel fine does not mean you can stop taking your medication. If you stop, serious complications of the disease may occur. Therefore, you should continue to take PRO-RAMIPRIL regularly, as prescribed by your doctor.

The "lifestyle" part of your treatment is as important as your medication. By working as a team with your doctor, you can help reduce the risk of complications to maintain the style of life you are accustomed to.

- **Alcohol:** Avoid alcoholic beverages until you have discussed their use with your doctor. Alcohol consumption may alter your blood pressure and/or increase the possibility of dizziness or fainting.
- **Diet:** Generally, avoid fatty foods and food that is high in salt or cholesterol.
- **Smoking:** Avoid it completely.

What it does:

PRO-RAMIPRIL is an angiotensin converting enzyme (ACE) inhibitor. You can recognize ACE inhibitors because their medicinal ingredient end in ‘-PRIL’.

This medicine does not cure your disease. It helps to control it. Therefore, it is important to continue taking PRO-RAMIPRIL regularly even if you feel fine.

When it should not be used:

Do not take PRO-RAMIPRIL if you:

- Are allergic to ramipril or to any non-medicinal ingredient in the formulation. Have experienced an allergic reactions (angioedema) with swelling of the hands, feet, or ankles, face, lips, tongue, throat, or sudden difficulty breathing or swallowing, to any ACE inhibitor or without a known cause. Be sure to tell your doctor, nurse, or pharmacist that this has happened to you.
- Have been diagnosed with hereditary angioedema: an increased risk of getting an allergic reaction that is passed down through families. This can be triggered by different factors, such as surgery, flu, or dental procedures.
- Are pregnant or intend to become pregnant. Taking PRO-RAMIPRIL during pregnancy can cause injury and even death to your baby.
- Are breastfeeding. PRO-RAMIPRIL passes into breast milk.
- Have narrowing of the arteries to one or both kidneys (renal artery stenosis).
- Have hypotension (low blood pressure).
- Are already taking a blood pressure-lowering medicine that contains aliskiren (such as Rasilez) and you have diabetes and/or kidney disease.

What the medicinal ingredient is:

ramipril.

What the important non-medicinal ingredients are:

Lactose monohydrate (spray-dried), magnesium stearate, talc and empty gelatin capsules (which are composed of titanium dioxide and/ or yellow iron oxide and / or FD & C red no. 40 and/or D & C red no. 28 and/or FD & C blue no. 1 and/ or D & C yellow no.10 and/or Black iron oxide)

What dosage forms it comes in:

Capsules 1.25 mg, 2.5 mg, 5.0 mg 10.0 mg and 15.0 mg.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions-Pregnancy

PRO-RAMIPRIL should not be used during pregnancy. If you discover that you are pregnant while taking PRO-RAMIPRIL, stop the medication and please contact your physician, nurse, or pharmacist as soon as possible.

BEFORE you use PRO-RAMIPRIL talk to your doctor, nurse or pharmacist if you:

- Are allergic to any drug used to lower blood pressure.
- Have recently received or are planning to get allergy shots for bee or wasp stings.
- Have narrowing of an artery or a heart valve.
- Have heart failure.
- Have diabetes, liver or kidney disease.
- Are on dialysis.
- Are dehydrated or suffer from excessive vomiting, diarrhea, or sweating.
- Are taking a salt substitute that contains potassium, potassium supplements, or potassium-sparing diuretic (a specific kind of “water pill”). Use of PRO-RAMIPRIL with these medicines is not recommended.
- Are on a low-salt diet.
- Are receiving gold (sodium aurothiomalate) injections.
- Are less than 18 years old.
- Are taking a medicine that contains aliskiren, such as Rasilez, used to lower high blood pressure. The combination with PRO-RAMIPRIL is not recommended.
- Are taking an angiotensin receptor blocker (ARB). You can recognize an ARB because its medicinal ingredient ends in “-SARTAN”.

You may become sensitive to the sun while taking PRO-RAMIPRIL. Exposure to sunlight should be minimized until you know how you respond.

If you are going to have surgery and will be given an anesthetic, be sure to tell your doctor or dentist that you are taking PRO-RAMIPRIL.

Driving and using machines: Before you perform tasks which may require special attention, wait until you know how you respond to PRO-RAMIPRIL. Dizziness, lightheadedness, or fainting can especially occur after the first dose and when the dose is increased.

Raynaud's phenomenon is a condition resulting from poor circulation in the extremities (i.e., fingers and toes). It may begin or get worse.

INTERACTIONS WITH THIS MEDICATION

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about all the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

The following may interact with PRO-RAMIPRIL

- Agents increasing serum potassium, such as a salt substitute that contains potassium, potassium supplements, or a potassium-sparing diuretic (a specific kind of “water pill”) or aliskiren. Use of

PRO-RAMIPRIL with these medicines is not recommended.

- Allopurinol used to treat gout.
- Antidiabetic drugs, including insulin and oral medicines.
- Lithium used to treat bipolar disease.
- Nonsteroidal anti-inflammatory drugs (NSAIDs), used to reduce pain and swelling. Examples include ibuprofen, naproxen, and celecoxib.
- Other blood pressure lowering drugs, including diuretics (“water pills”). When taken in combination with PRO-RAMIPRIL, they may cause excessively low blood pressure.
- Nitrates
- ASA (aspirin)
- Heparin
- Immunosuppressants
- Corticosteroids
- Procainamide
- Cytostatics
- Other substances that may change the normal results expected to be measured on a routine blood test.
- Blood pressure-lowering drugs, including diuretics (“water pills”), aliskiren-containing products (e.g. Rasilez), or angiotensin receptor blockers (ARBs).

PROPER USE OF THIS MEDICATION

Take PRO-RAMIPRIL exactly as prescribed. It is recommended to take your dose at about the same time everyday.

Usual adult dose:

High Blood Pressure: The recommended initial dosage of PRO-RAMIPRIL is 2.5 mg once daily. Your doctor will determine the appropriate dosage.

For patients taking diuretics (“water pills”) or with impaired kidney function: The recommended initial dosage of PRO-RAMIPRIL is 1.25 mg daily.

Overdose:

If you think you have taken too much PRO-RAMIPRIL contact your doctor, nurse, pharmacist, hospital emergency department or regional Poison control Centre immediately, even if there are no symptoms.

Missed Dose:

If you forgot to take your dose during the day, carry on with the next one at the usual time. Do not double dose.

IMPORTANT: PLEASE READ

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- Dizziness
- Drowsiness, fatigue, weakness
- Cough
- Rash, itching, flushing
- Headache
- Abdominal pain
- Sad mood

If any of these affects you severely, tell your doctor, nurse or pharmacist.

Other side effects may include: difficulty in maintaining your balance while standing, nasal or sinus congestion, swollen lymph nodes, bronchitis, loss of hair, impotence/reduced libido, difficulty with sleep, restlessness, inflammation of the eye (pink eye), taste modifications or loss of taste, vision or hearing modifications, attention disturbances, skin inflammation or red skin, burning sensation, inflammation of the mouth or tongue, aggravated asthma, breast enlargement in males.

PRO-RAMIPRIL can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Common	Low Blood Pressure: dizziness, fainting, lightheadedness	✓		
	Increased levels of potassium in the blood: irregular heartbeat, muscle weakness and generally feeling unwell		✓	
Uncommon	Allergic Reaction: rash, hives, swelling of the face, arms and legs, lips, tongue or throat, difficulty swallowing or breathing			✓

	Kidney Disorder: change in frequency of urination, nausea, vomiting, swelling of extremities, fatigue		✓	
	Liver Disorder: yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		✓	
	Electrolyte Imbalance: weakness, drowsiness, muscle pain or cramps, irregular heartbeat		✓	
Rare	Decreased Platelets: bruising, bleeding, fatigue, and weakness		✓	
	Decreased White Blood Cells: infections, fatigue, fever, aches, pains, and flu-like symptoms		✓	
	Heart Attacks: chest pain and/or discomfort, pain in the jaw, shoulders, arm and/or back, shortness of breath, sweating, lightheadedness, nausea			✓
	Cerebro-vascular accidents/Stroke: weakness, trouble speaking, trouble seeing, headaches, dizziness			✓
	Intestinal Angiodema: abdominal pain (with or without nausea or vomiting)			✓

This is not a complete list of side effects. For any unexpected effects while taking PRO-RAMIPRIL, contact your doctor or pharmacist.

HOW TO STORE IT

Store at room temperature 15-30°C, (59-86°F) in a well-closed container.

Keep this medication out of the reach and sight of children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator: 0701E
Ottawa, Ontario
K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

Note: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

For more information, please contact your doctor, pharmacist or other healthcare professional.

This document plus the full product monograph, prepared for health professionals, can be obtained by contacting Pro Doc Ltée at 1-800-361-8559, www.prodoc.qc.ca or info@prodoc.qc.ca.

This leaflet was prepared by
Pro Doc Ltée, Laval, Québec, H7L 3W9

Date of Revision: October 22, 2014