

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

PrCEPHALEXIN - 500

Cephalexin Tablets USP

500 mg

Read this carefully before you start taking CEPHALEXIN - 500 and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about CEPHALEXIN - 500.

What is CEPHALEXIN - 500 used for?

- CEPHALEXIN - 500 is used to treat certain bacterial infections in the:
 - respiratory tract;
 - ear (otitis media);
 - genitals and urinary tract;
 - bones and joints;
 - skin and soft tissue.
- Antibacterial drugs like CEPHALEXIN - 500 treat only bacterial infections. They do not treat viral infections, such as the common cold.

How does CEPHALEXIN - 500 work?

CEPHALEXIN - 500 is an antibiotic that:

- Stops the growth of bacteria
- Kills bacteria

What are the ingredients in CEPHALEXIN - 500?

Medicinal ingredients: Cephalexin

Non-medicinal ingredients: Colloidal silicon dioxide, FD&C yellow #6, hydroxypropyl cellulose, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, stearic acid, and titanium dioxide.

CEPHALEXIN - 500 comes in the following dosage forms:

500 mg Tablets: Supplied in bottles of 100 and 500 Tablets.

Do not use CEPHALEXIN - 500 if:

- you are allergic to cephalexin, cephalosporin antibiotics or to any of the other ingredients in CEPHALEXIN - 500

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take CEPHALEXIN - 500. Talk about any health conditions or problems you may have, including if you:

- you are allergic to or react badly to penicillins or other antibiotics.
- you have intestinal or bowel problems.
- you have kidney problems.
- you are pregnant or plan to be pregnant.
- you are breast-feeding or plan to breast-feed.

Other warnings that you should know about:

- Using antibiotics like CEPHALEXIN - 500 may cause with *Clostridium difficile*-associated disease (CDAD). See “Serious side effects and what to do about them”, below.
- CEPHALEXIN - 500 may interfere with some blood and urine test results. Talk to your doctor if you are given a blood or urine test while taking CEPHALEXIN - 500.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

How to take CEPHALEXIN - 500:

- Take CEPHALEXIN - 500 orally on an empty stomach.
- Although you may feel better early in treatment, CEPHALEXIN - 500 should be used exactly as directed.
- Misuse or overuse of CEPHALEXIN - 500 could lead to the growth of bacteria that will not be killed by CEPHALEXIN - 500 (resistance). This means that CEPHALEXIN - 500 may not work for you in the future.
- Do not share your medicine.

Usual dose:

Adult:

The adult dosage ranges from 1 to 4 g daily in divided doses. The usual adult dose is 1 g/day in divided doses every 6 hours.

Children:

The recommended daily dosage for children is 25 to 50 mg/kg/day in divided doses every 6 hours.

Overdose:

Symptoms of oral overdose may include:

- nausea
- vomiting
- abdominal pain
- diarrhea
- bloody urine

If you think you have taken too much CEPHALEXIN - 500, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

What are possible side effects from using CEPHALEXIN - 500?

These are not all the possible side effects you may feel when taking CEPHALEXIN - 500. If you experience any side effects not listed here, contact your healthcare professional.

You may experience diarrhea, nausea, vomiting, loose stools, abdominal pain, diaper rash, inflammation of the vagina or discharge, dizziness, fatigue, headache, agitation, confusion, hallucinations or symptoms that you do not understand. You should tell your health professional for of any of these symptoms as soon as possible.

Vertigo (loss of balance or unsteadiness), tinnitus (ringing in the ears), hearing loss and behavioural changes in young children have been reported.

CEPHALEXIN - 500 may also cause effects such as yellowing of the whites of the eyes or skin (jaundice) or the inflammation of the liver (hepatitis).

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE			
Symptoms of a severe bowel condition (<i>Clostridium difficile</i> colitis): <ul style="list-style-type: none"> • persistent diarrhea • bloody or watery diarrhea • abdominal or stomach pain/cramping • blood/mucus in stool 			√
Severe skin reactions: <ul style="list-style-type: none"> • skin rash, which may blister, and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge) • widespread rash with blisters and skin peeling on much of the body surface particularly around the mouth, nose, eyes and genitals. 			√
Severe Cutaneous Adverse Reactions (SCAR) (severe skin reactions that may also affect other organs): <ul style="list-style-type: none"> • Skin peeling, scaling, or blistering (with or without pus) which may also affect your eyes, mouth, nose or genitals, itching, severe rash, bumps under the skin, skin pain, skin color changes (redness, yellowing, purplish) • Swelling and redness of eyes or face • Flu-like feeling, fever, chills, body aches, swollen glands, cough Shortness of breath, chest pain or discomfort			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Tablets:

Store at room temperature (15°C to 30°C) in a tightly closed container. Dispense in a light-resistant container, as prolonged exposure to light may cause fading of the coating.

If you want more information about CEPHALEXIN - 500:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); or by contacting Pro Doc Ltée at 1-800-361-8559, www.prodoc.qc.ca or info@prodoc.qc.ca.

This leaflet was prepared by
Pro Doc Ltée, Laval, Québec, H7L 3W9

Last revised: May 28, 2021